

Isovaleric Acidemia: New Aspects of Genetic and Phenotypic Heterogeneity

JERRY VOCKLEY* AND REGINA ENSENAUER

Isovaleric acidemia (IVA) is an autosomal recessive inborn error of leucine metabolism caused by a deficiency of the mitochondrial enzyme isovaleryl-CoA dehydrogenase (IVD) resulting in the accumulation of derivatives of isovaleryl-CoA. It was the first organic acidemia recognized in humans and can cause significant morbidity and mortality. Early diagnosis and treatment with a protein restricted diet and supplementation with carnitine and glycine are effective in promoting normal development in severely affected individuals. Both intra- and interfamilial variability have been recognized. Initially, two phenotypes with either an acute neonatal or a chronic intermittent presentation were described. More recently, a third group of individuals with mild biochemical abnormalities who can be asymptomatic have been identified through newborn screening of blood spots by tandem mass spectrometry. IVD is a flavoenzyme that catalyzes the conversion of isovaleryl-CoA to 3-methylcrotonyl-CoA and transfers electrons to the electron transfer flavoprotein. Human IVD has been purified from tissue and recombinant sources and its biochemical and physical properties have been extensively studied. Molecular analysis of the *IVD* gene from patients with IVA has allowed characterization of different types of mutations in this gene. One missense mutation, 932C>T (A282V), is particularly common in patients identified through newborn screening with mild metabolite elevations and who have remained asymptomatic to date. This mutation leads to a partially active enzyme with altered catalytic properties; however, its effects on clinical outcome and the necessity of therapy are still unknown. A better understanding of the heterogeneity of this disease and the relevance of genotype/phenotype correlations to clinical management of patients are among the challenges remaining in the study of this disorder in the coming years. © 2006 Wiley-Liss, Inc.

KEY WORDS: isovaleric acidemia; organic acidemia; flavoenzyme; leucine metabolism; acyl-CoA dehydrogenase

INTRODUCTION

Isovaleric acidemia (IVA) was the first of the organic acidemias to be described and its nature was elucidated due to a combination of astute clinical acumen and new technology. The two original patients were 4 and 2½ years old, respectively, and had similar histories of recurrent episodes of vomiting and lethargy that resolved with supportive therapy including glucose infusions [Tanaka et al., 1966; Budd et al., 1967; Efron, 1967]. A “smell specialist” sug-

gested that the unusual odor associated with their acute episodes was likely a short chain fatty acid, and isovaleric acid was identified in patient plasma through the then novel approach of gas chromatography followed by mass spectrometry [Tanaka, 1990]. Urine was subsequently shown to contain isovalerylglycine and 3-hydroxyisovaleric acid as well as other metabolites [Tanaka and Isselbacher, 1967; Tanaka et al., 1968]. Oxidation of labeled isovaleric acid by patient white blood cells was reduced compared to

control cells, and a defect of leucine metabolism was postulated [Shih et al., 1973; Rhead and Tanaka, 1980; Yoshida et al., 1985; Hyman and Tanaka, 1986]. It must be remembered that until the characterization of these patients, the oxidation of isovaleryl-CoA was assumed to be catalyzed by the newly identified short chain acyl-CoA dehydrogenase [Crane et al., 1956; Engel and Massey, 1971; McKean et al., 1979; Frerman et al., 1980]. Over the next 20 years IVD was separated from the other acyl-CoA dehydrogenases, purified, cloned, and the molecular nature of the defects responsible for loss of IVD activity elucidated. The use of gas chromatography/mass spectrometry (GC-MS) ultimately became the mainstay of the identification and routine clinical diagnosis of a new class of inborn errors of metabolism resulting in the abnormal accumulation of organic acids in urine, and it remains a valuable tool for biochemical geneticists today [Tanaka et al., 1980; Burke et al., 1983]. The

Jerry Vockley is Professor of Pediatrics and Human Genetics at the University of Pittsburgh School of Medicine and Graduate School of Public Health, and Chief of Medical Genetics at the Children's Hospital of Pittsburgh. He has extensively researched the molecular and biochemical basis of isovaleric acidemia.

Regina Ensenauer is a pediatrician and biochemical geneticist at the Children's Hospital of the Ludwig-Maximilians University in Munich, Germany. She currently holds an academic career research scholarship. Her research interests are in the field of branched-chain amino acid catabolism and fatty acid oxidation.

Grant sponsor: NIH; Grant number: RO1-DK45482; Grant sponsor: Pennsylvania Tobacco Settlement Fund.

*Correspondence to: Jerry Vockley, Department of Pediatrics, University of Pittsburgh School of Medicine, The Children's Hospital of Pittsburgh, 3705 Fifth Avenue, Pittsburgh, PA 15238.

E-mail: gerard.vockley@chp.edu

DOI 10.1002/ajmg.c.30089

Published online 00 Month 2006 in Wiley InterScience (www.interscience.wiley.com)

more recent application of tandem mass spectrometry (MS/MS) for analysis of the acylcarnitine profile of blood spots from newborn screening filter paper cards has allowed a significant expansion of the recognition of mildly affected and potentially asymptomatic individuals with IVD deficiency through newborn screening [Ensenauer et al., 2004].

DISEASE SPECTRUM

Early literature on IVA, an autosomal recessive disorder, emphasized two apparent phenotypes [Tanaka, 1990]. The first was an acute, neonatal presentation with patients becoming symptomatic within the first 2 weeks of life [Tanaka et al., 1966; Budd et al., 1967; Efron, 1967; Lott et al., 1972;

Early literature on IVA, an autosomal recessive disorder, emphasized two apparent phenotypes. The first was an acute, neonatal presentation with patients becoming symptomatic within the first 2 weeks of life.

Levy et al., 1973; Elsas and Naglak, 1988]. Patients appeared initially well, then developed vomiting and lethargy, progressing to coma. The second group presented with relatively non-specific failure to thrive and/or developmental delay (chronic intermittent presentation) [Levy et al., 1973; Shih et al., 1984; Berry et al., 1988; Elsas and

The second group presented with relatively non-specific failure to thrive and/or developmental delay.

Naglak, 1988; Mehta et al., 1996]. Patients who survived an early acute presentation subsequently were indis-

tinguishable from those with the chronic phenotype, and both groups of patients were prone to intermittent acute episodes of decompensation with minor illnesses [Tanaka, 1990]. In reality it is now apparent that patients can fall anywhere on the spectrum of acute to chronic presentation and that there is probably little predictive value to the initial presentation. Moreover, with the application of MS/MS in newborn screening, potentially asymptomatic patients with one recurring IVD gene mutation and a mild biochemical phenotype are being identified in increasing numbers, representing an additional phenotype of IVA [Ensenauer et al., 2004]. This type may represent a biochemical phenotype only without the expression of any clinical symptoms (such as in benign hyperphenylalaninemia) and, therefore, needs to be differentiated from the classic forms of IVA. For practical purposes, we suggest classifying patients with IVA as “metabolically severe” and “metabolically mild or intermediate”, giving consideration to the broader spectrum of IVA detectable by newborn screening.

CLINICAL PRESENTATION

Nearly all published clinical information on patients with IVA is retrospective [Sweetman and Williams, 2001], and thus the following discussion of clinical symptoms is limited to the classic presentations of IVA prior to newborn screening, that is, manifestation in the neonatal period versus later in childhood. In one summary of 37 patients compiled from different publications, 28 presented in the first 2 weeks of life, 7 between 2 weeks and 1 year of age, and the remaining 2 after 1 year of age [Tanaka, 1990]. Sixteen of the patients were deceased, and of those still alive, seven were reported to have mild to moderate mental retardation.

Neonatal symptoms are non-specific and include poor feeding, vomiting, decreased level of consciousness, and seizures [Tanaka et al., 1966; Budd et al., 1967; Efron, 1967; Lott et al., 1972; Levy et al., 1973; Spirer et al., 1975; Elsas and Naglak, 1988]. Infants may develop hypothermia and appear to be dehy-

drated. A characteristic smell of “dirty socks” may be present when the patient is acutely sick though, unlike other organic acidemias, the urine has no odor since the unconjugated isovaleric acid responsible for the odor is not excreted in urine in appreciable quantity [Tanaka et al., 1966; Tanaka, 1990]. The odor

A characteristic smell of “dirty socks” may be present when the patient is acutely sick though, unlike other organic acidemias, the urine has no odor since the unconjugated isovaleric acid responsible for the odor is not excreted in urine in appreciable quantity. The odor may be best appreciated in body sweat or cerumen from the ear.

may be best appreciated in body sweat or cerumen from the ear. Acidosis with an unexplained anion gap is characteristic, and hyperammonemia, hyper- or hypoglycemia and hypocalcemia may be present [Tanaka et al., 1966; Budd et al., 1967; Lott et al., 1972; Levy et al., 1973; Yoshino et al., 1982; Mendiola et al., 1984; Tanaka, 1990; Worthen et al., 1994]. Secondary hyperammonemia is presumed to be due to inhibition of N-acetylglutamate synthetase by isovaleryl-CoA and/or intracellular depletion of acetyl-CoA leading to reduced N-acetylglutamate synthesis and impairment of the urea cycle [Coude et al., 1979; Stewart and Walser, 1980]. Pancytopenia as well as isolated neutropenia and thrombocytopenia can occur due to bone marrow suppression [Kelleher et al., 1980]. Left untreated, patients may progress to coma and death often due to cerebral edema or hemorrhage [Fischer et al., 1981]. Overall, the clinical picture overlaps other organic acidemias including the β -oxidation defects as well as the urea cycle disorders and other primary causes of hyperammonemia, all of which

must, therefore, be considered in the differential diagnosis. Patients who survive a neonatal crisis may be clinically indistinguishable from children diagnosed later in life [Tanaka, 1990].

Children diagnosed outside the newborn period may present with more chronic, relatively non-specific findings of failure to thrive and/or developmental delay or mental retardation [Tanaka, 1990]. Minus the “sweaty feet odor” of isovaleric acid, which is not present when a patient is otherwise well, there is little to suggest a specific diagnosis in these children and thus, it must be considered in all patients with this clinical picture. They also are at risk of episodes of acute acidosis and metabolic decompensation, usually due to intercurrent illnesses or other physiologic stress including fasting [Berry et al., 1988; Tanaka, 1990]. Acute episodes may be misdiagnosed as diabetic ketoacidosis due to hyperglycemia, acidosis and the apparent presence of blood and urinary ketones [Attia et al., 1996]. Acute pancreatitis, myeloproliferative syndrome, Fanconi syndrome, and cardiac arrhythmias have been reported [Arnold et al., 1986; Kahler et al., 1994; Weinberg et al., 1997; Gilbert-Barnes and Barnes, 1999]; abnormalities of the globus pallidus can be seen [Sogut et al., 2004]. The variability of this disorder is highlighted by the diagnosis of IVA in a previously well 18-year-old man who developed acute nausea, vomiting, and mental status changes during basic training camp for the United States Air Force [Feinstein and O'Brien, 2003].

There have been reports of successful pregnancies in women with IVA resulting in apparently well infants [Shih et al., 1984; Spinty et al., 2002].

BIOCHEMICAL DIAGNOSIS AND FOLLOW-UP

The majority of patients with IVA today are diagnosed pre-symptomatically through newborn screening by use of MS/MS which reveals elevations of the marker metabolite C5 acylcarnitine in dried blood spots. Because C5 acylcarnitine represents a mixture of isomers

(isovalerylcarnitine, 2-methylbutyrylcarnitine, and pivaloylcarnitine), further diagnostic evaluation is required (Table I). Elevations of 2-methylbutyrylcarnitine are seen in patients with 2-methylbutyrylglycinuria caused by a deficiency of short/branched-chain acyl-CoA dehydrogenase (SBCAD), an inborn error of isoleucine catabolism [Andresen et al., 2000; Gibson et al., 2000], whereas pivaloylcarnitine is derived from pivalic acid, a component of several antibiotics [Abdenur et al., 1998]. Isovaleryl-CoA intermediates can also be seen in deficiencies of the electron transfer flavoprotein (ETF) and its dehydrogenase (glutaric aciduria type 2).

A long list of other isovaleryl-CoA derived metabolites has been reported in blood and urine from patients with IVA and can assist in confirmation of the disorder [Lehnert, 1981a,b, 1983; Lehnert and Niederhoff, 1981; Burke et al., 1983; Dorland et al., 1983; Hine and Tanaka, 1984; Poorthuis et al., 1993; Loots et al., 2005]. Isovaleryl conjugates of multiple amino acids have also been detected in urine, as have free 3- and 4-hydroxyisovaleric acids [Tanaka et al., 1968; Lehnert and Niederhoff, 1981; Loots et al., 2005]. Free isovaleric acid in blood during episodes of acute metabolic decompensation can reach several hundred times normal values but is not readily seen in blood and urine due to its rapid conjugation to other compounds. Thus, isovalerylcarnitine and isovalerylglycine are the hallmarks of this

disorder in plasma and urine, respectively, and are elevated regardless of a patient's metabolic condition. Quantification of both conjugates has suggested

Free isovaleric acid in blood during episodes of acute metabolic decompensation can reach several hundred times normal values but is not readily seen in blood and urine due to its rapid conjugation to other compounds. Thus, isovalerylcarnitine and isovalerylglycine are the hallmarks of this disorder in plasma and urine, respectively, and are elevated regardless of a patient's metabolic condition.

a correlation of the metabolite concentrations with genotype, differentiating between groups of patients with a metabolically severe phenotype associated with heterogeneous *IVD* gene mutations and patients with a metabolically mild or intermediate phenotype associated with one recurring mutation [Ensenauer et al., 2004]. Disease-specific

TABLE I. Initial Diagnostic Evaluation for Isovaleric Acidemia

Evaluation following abnormal newborn screening with an elevated C5 acylcarnitine concentration	
Test	Determination of
Urine organic acid analysis	Multiple abnormal metabolites; isovalerylglycine concentration
Plasma acylcarnitine analysis	Isovalerylcarnitine concentration
Plasma carnitine analysis	Free carnitine concentration
Molecular genetic analysis	Common 932C>T (A282V) <i>IVD</i> gene mutation associated with a mild biochemical phenotype; otherwise heterogeneous mutations
Enzymatic analysis optional (fibroblasts, lymphocytes)	Residual enzyme activity

metabolites also accumulate in amniotic fluid during pregnancy with an affected fetus and provide the opportunity for prenatal diagnosis [Jakobs et al., 1984; Hine et al., 1986; Shigematsu et al., 1991].

Several direct and indirect methods to assay IVD activity have been published and, in addition to molecular genetic analysis, can be used to confirm a diagnosis of IVA [Shih et al., 1973; Rhead and Tanaka, 1980; Yoshida et al., 1985; Hyman and Tanaka, 1986; Vockley et al., 1991; Mohsen et al., 1998; Tajima et al., 2005]. Fibroblasts, lymphocytes, and amniocytes all have measurable amounts of IVD activity and serve as ready sources of tissue for this purpose [Vockley et al., 1991; Kleij et al., 1995; Mohsen et al., 1998; Ensenaer et al., 2004]. While significant residual activity blunts the level of abnormal metabolites, correlation between clinical presentation and enzyme activity has been poor [Ikeda et al., 1985b; Hyman and Tanaka, 1986; Vockley et al., 1991].

Regarding routine follow-up visits, there is no established laboratory marker for monitoring therapeutic control or disease state. Weight gain, growth and development should be age-appropriate and thus, body measurements are key parameters to follow on a routine basis. Specifically, protein malnutrition must be avoided if the patient is protein


restricted. Analysis of amino acids, albumin, and prealbumin in plasma is recommended to monitor this. Plasma levels of leucine are not elevated in IVA, even if untreated, due to the irreversible oxidative decarboxylation earlier in the leucine degradation pathway [Sweetman and Williams, 2001]. Plasma free carnitine concentrations may be helpful for determining necessity and monitoring of carnitine supplementation.

MOLECULAR FINDINGS

IVD is an intramitochondrial homotetrameric flavoenzyme that catalyzes the α,β -dehydrogenation of isovaleryl-CoA resulting in 3-methylcrotonyl-CoA (Fig. 1) and transfers the reducing equivalents to the ETF [Ikeda et al., 1983; Ikeda and Tanaka, 1988; Mohsen and Vockley, 1995]. It is encoded in the nuclear genome as a precursor protein, but is active in the mitochondrial matrix [Vockley et al., 1991; Ikeda et al., 1987; Parimoo and Tanaka, 1993]. Following transcription in the nucleus and translation in the cytoplasm, it is held in a partially folded state by chaperonin proteins, targeted to mitochondria by a characteristic amino terminal peptide sequence and imported into mitochondria. The target peptide is cleaved in the mitochondrial matrix, folding of the monomer is completed,

and the final active tetramer is assembled. The FAD cofactor is probably added after import into the mitochondria, but the exact timing and mechanism of this reaction is poorly characterized [Nagao and Tanaka, 1992]. The reaction catalyzed by IVD is initiated upon acyl-CoA substrate binding. In the reductive half-reaction, the formation of a hydrogen bond between the acyl carbonyl oxygen and the 2'-hydroxyl hydrogen of the FAD ribityl moiety is crucial for activation of the acyl moiety^{Q1} [Ghisla et al., 1992; Nishina et al., 1995; Miura et al., 1996]. The C α and C β hydrogens of the acyl moiety are then removed in concert as a proton and a hydride, respectively, forming a stable charge-transfer complex. The ETF, considered a second substrate, extracts the reducing equivalents from the charge-transfer complex in the oxidative half-reaction during which electrons are transferred to the latter and the enoyl-CoA product is released [Gorelick et al., 1985; Ikeda et al., 1985a; Ghisla and Massey, 1989]. The molecular mechanism of the oxidative half-reaction and the order of binding of the substrates have not been elucidated.

Molecular cloning of the *IVD* gene showed it to be located on chromosome 15q14-15, consisting of 12 exons that span ~15 kb of genomic DNA [Parimoo and Tanaka, 1993]. Molecular analysis of the *IVD* gene in patients with



symptomatic IVA has identified numerous point mutations in the protein coding region that lead to the production of an inactive or unstable protein (Fig. 2) [Ikeda et al., 1985b; Vockley et al., 1991, 1992a,b, 2000; Mohsen et al., 1998]. Some of these have been shown to cause only a mild alteration in enzyme function, correlating to some extent with a mild clinical phenotype in patients [Mohsen et al., 1998; Nasser et al., 2004]. A significant proportion of the mutant alleles lead to abnormal splicing of the IVD RNA and subsequent complete lack of IVD protein [Vockley et al., 1991, 2000]. Studies of the bioprocessing of IVD protein in fibroblasts from patients with IVA have reflected the effects of these mutations but have not provided insight into the clinical variability seen in the disorder [Ikeda et al., 1985b; Vockley et al., 1991]. In general, genotype and phenotype have not been well correlated.

NEWBORN SCREENING AND ISOVALERIC ACIDEMIA

With the advent of the use of MS/MS to screen newborn blood spots for acylcarnitine concentrations, most if not all patients with IVA should be identified as newborns prior to the development of symptoms. One unexpected finding to

arise from newborn screening studies is the identification of individuals with only mild elevations of isovaleryl-CoA related metabolites in plasma and urine, orders of magnitude lower than in the classic forms of IVA, and apparently only partial reduction in IVD activity (Table II) [Ensenauer et al., 2004]. Nearly half of the mutant IVD alleles sequenced from infants diagnosed by newborn screening have been found to contain a common recurring missense mutation [Ensenauer et al., 2005] (932C>T; A282V; Fig. 2). This specific mutation was present in approximately two thirds of newborns in this cohort, mostly in a compound heterozygous fashion. All of the affected newborns carrying the common mutation have remained asymptomatic with mild or no dietary protein restriction and carnitine supplementation if necessary over a maximum duration of follow-up of up to 5 years of age. Subsequently, asymptomatic siblings of patients identified through newborn screening (ages 3–11 years at the time of diagnosis) have been found to be homozygous or compound heterozygous for the same mutation with a similarly mild biochemical phenotype [Ensenauer et al., 2004]. They have remained without symptoms during episodes of febrile illnesses.

Prior to newborn screening, this mutation was identified only in a single

patient with mild IVD deficiency originally evaluated for a tic disorder and slight developmental delay [Vockley et al., 1992b; Mohsen et al., 1998]. The mutant A282V protein is stable in vitro but is kinetically impaired, exhibiting an increased K_m and a reduced catalytic efficiency, and has diminished thermal stability [Mohsen et al., 1998; Nasser et al., 2004]. It is clear that the newborn screening patients who carry the common mutation either in a homozygous or compound heterozygous state and their sibs skew the spectrum of IVA with more than half of individuals representing a new mild phenotype and potentially remaining asymptomatic. This is an expansion

It is clear that the newborn screening patients who carry the common mutation either in a homozygous or compound heterozygous state and their sibs skew the spectrum of IVA with more than half of individuals representing a new mild phenotype and potentially remaining asymptomatic.




Figure 2. Molecular defects in the *IVD* gene on chromosome 15q14–15 [Mohsen et al., 1998; Ensenauer et al., 2004]. The nucleotide change 932C>T (A282V) in exon 9 has been identified in a significant proportion of individuals diagnosed by newborn screening [Ensenauer et al., 2004].

TABLE II. Recommendations for Therapy in Isovaleric Acidemia

Therapy	Biochemical phenotype	
	Metabolically mild or intermediate ^a	Metabolically severe ^b
Prevention of metabolic crisis	Close clinical observation; promote anabolism during illness	
Diet	None	Protein restriction
Medication	Carnitine (30–50 mg/kg per day) if plasma free carnitine concentration is low	Carnitine (100 mg/kg per day)
	None	Glycine (150–250 mg/kg per day)

^aMetabolically mild or intermediate: NBS blood spot C5 acylcarnitine concentration 0.8–6 $\mu\text{mol/L}$; urine isovalerylglycine concentration 15–195 mmol/mol creatinine.

^bMetabolically severe: NBS blood spot C5 acylcarnitine concentration up to 21.7 $\mu\text{mol/L}$; urine isovalerylglycine concentration up to 3,300 mmol/mol creatinine.

Values derived from Ensenauer et al., 2004.

of our view of the natural history of IVA prior to the newborn screening era and leads to significant implications for management and genetic counseling. As opposed to the classic forms of IVA, it is still uncertain whether individuals with the latter type have a disease, a risk of clinical manifestation, or simply express a clinically insignificant biochemical phenotype. While these individuals may have normal leucine homeostasis under physiological conditions, their risk of metabolic decompensation under stress conditions remains to be elucidated.

THERAPY

There are three goals for therapy of IVA (Table II). The first is prevention of metabolic decompensation by careful clinical observation of the patient regardless of the type of IVA. During times of metabolic stress (including illness and fasting) endogenous leucine from protein catabolism adds significantly to the production of isovaleryl-CoA [Collins et al., 1987; Millington et al., 1987; Pollitt, 1987]. Achieving anabolism is the main therapeutic approach. Thus, sick day precautions for patients with IVA should include increased caloric intake in addition to decreased leucine intake. This is most easily accomplished with oral solutions containing simple sugars and leucine free metabolic formulae or powders. IV glu-

cose infusions need to be added if oral intake is interrupted. Leucine intake should be decreased to approximately 50% of the patient's usual daily minimum, but returned to normal after 24 hr in order to promote protein anabolism.

The second goal is long-term reduction of the production of isovaleryl-CoA from leucine catabolism through dietary manipulation [Lott et al., 1972; Levy et al., 1973; Berry et al., 1988; Sweetman and Williams, 2001]. Total protein and caloric intake must be adequate to support normal growth in children and maintain an anabolic state, and thus monitoring of weight, length, and head circumference is essential at follow-up. In many cases, it may be sufficient to moderately lower protein intake with natural foods to approximately 1.5 gm/kg per day. In patients with recurrent clinical symptoms, leucine restriction in excess of total natural protein may also be necessary [Sweetman and Williams, 2001]. Natural protein necessary to reach the recommended age-appropriate daily requirement must then be provided with leucine-free amino acids. Because of the specific role of leucine in promoting protein synthesis, however, there is a potential for adverse side effects of rigorous leucine restriction including muscle wasting [Harris et al., 2004].

Acute episodes of metabolic decompensation can present with emesis, lethargy and signs of overwhelming

acidosis. Under these circumstances, immediate hospitalization is required so that IV access can be established and glucose administered. Glucose infusion should be calculated to give at least 8 mg/kg per min with concomitant use of IV insulin if necessary to maintain euglycemia. Reintroduction of oral intake including catabolism-sparing levels of protein (0.5 gm/kg per day) with leucine should occur as soon as it can be tolerated, otherwise parenteral amino acids should be provided. If present, hyperammonemia will reverse with correction of the primary metabolic derangement; alternative ammonia conjugating agents such as sodium benzoate or phenylbutyrate are generally not indicated.

The third goal of therapy in patients with IVA is to prevent the accumulation of toxic metabolites by shunting isovaleryl-CoA towards reactions that produce metabolites presumed to be non-toxic and that can readily be excreted. Recognition of isovalerylglycine in urine in the initial patients with IVA first led to the use of glycine to achieve this end [Krieger and Tanaka, 1976; Cohn et al., 1978; Yudkoff et al., 1978; Elsas and Naglak, 1988]. Isovaleryl-CoA is enzymatically conjugated to glycine, a reaction that can be augmented by supplementation with exogenous glycine to supra-physiologic levels. Such supplementation prevents or reduces the accumulation of isovaleric

acid in blood following a leucine load, and the length and severity of symptoms during intercurrent illnesses [Krieger and Tanaka, 1976; Yudkoff et al., 1978; Shigematsu et al., 1982]. Doses of 150–

Isovaleryl-CoA is enzymatically conjugated to glycine, a reaction that can be augmented by supplementation with exogenous glycine to supra-physiologic levels. Such supplementation prevents or reduces the accumulation of isovaleric acid in blood following a leucine load, and the length and severity of symptoms during intercurrent illnesses.

600 mg/kg per day given orally and divided in three or four equal doses of body weight have been proposed, but the optimum dose has not been determined. Patients exhibit a dose sensitive increase in excretion of isovalerylglycine, but at least in one report an increase in the glycine dose from 300 to 600 mg/kg of body weight led to a decrease in the excretion of isovalerylglycine, presumably due to inhibition of glycine-N-acylase by glycine [Elsas and Naglak, 1988]. Thus, initial dosing in the range of 150–250 mg/kg per day is reasonable in patients with a metabolically severe type of IVA (Table II). Concern has been raised about the potential for glycine toxicity, though no reports of such an occurrence have been published.

The identification of isovalerylcarnitine in blood and urine along with the frequent observation of a secondary deficiency of free carnitine in patients with IVA has prompted treatment with carnitine. A dose of 100 mg/kg body weight per day has generally been used (Table II) and has been shown to increase the excretion of isovalerylcarnitine in urine [Mayatepek et al., 1991; Fries

et al., 1996]. Combined therapy with carnitine and glycine has been shown to maximize the total excretion of isovaleryl-CoA conjugates, but the clinical benefit of combined versus single therapy has not been established through controlled studies [Fries et al., 1996; Itoh et al., 1996]. The relative merits of the two therapies either singly or together in patients with more severe presentations including recurrent crises remains a matter of debate.

The necessity of any treatment for individuals diagnosed by newborn screening and carrying the common 932C>T (A282V) mutation is unclear. Specifically, the potential for metabolic decompensation under stress conditions remains to be elucidated. It appears reasonable to observe these individuals clinically, particularly when exposed to metabolic stressors such as febrile illnesses or fasting (e.g., when undergoing surgery). Additional recommendations include low-dose carnitine supplementation if the plasma free carnitine concentration is reduced (Table II).

SUMMARY

IVA was originally viewed as a relatively rare, life threatening inborn error of metabolism with both acute and chronic manifestations. Recent data from newborn screening studies and additional molecular and cellular laboratory investigations have revealed a far more heterogeneous condition with a potential for normal growth and development. Prospective long-term follow-up of newborns identified with IVA and clinical trials of carnitine and glycine therapy will be critical to optimization of outcome in these patients.

ACKNOWLEDGMENTS

This review is dedicated to the memory of Professor Kay Tanaka who originally identified isovaleric acidemia. Dr. Tanaka died on August 21, 2005 at the age of 75. JV was supported in part by NIH grant RO1-DK45482 and the Pennsylvania Tobacco Settlement Fund. RE was supported by the “Hochschul- und Wissenschaftsprogramm” of the

Ludwig-Maximilians University Munich, Germany.

REFERENCES

- Abdenur JE, Chamoles NA, Guinle AE, Schenone AB, Fuertes AN. 1998. Diagnosis of isovaleric acidemia by tandem mass spectrometry: False positive result due to pivaloylcarnitine in a newborn screening programme. *J Inher Metab Dis* 21: 624–630.
- Andresen B, Christensen E, Corydon T, Bross P, Pilgaard B, Wanders R, Ruitter J, Simonsen H, Winter V, Knudsen I, Schroeder LD, Gregersen N, Skovby F. 2000. Isolated 2-methylbutyrylglycinuria caused short/branched-chain acyl-CoA dehydrogenase deficiency: Identification of a new enzyme defect, resolution of its molecular basis, and evidence for distinct acyl-CoA dehydrogenases in isoleucine and valine metabolism. *Am J Hum Gen* 67:1095–1103.
- Arnold WC, Brewster M, Byrne WJ, Booth B. 1986. Fanconi syndrome in a patient with a variant of isovaleric acidemia. *Intl J Pediatr Nephrol* 7:95–98.
- Attia N, Sakati N, al Ashwal A, al Saif R, Rashed M, Ozand PT. 1996. Isovaleric acidemia appearing as diabetic ketoacidosis. *J Inherited Metab Disease* 19:85–86.
- Berry GT, Yudkoff M, Segal S. 1988. Isovaleric acidemia: Medical and neurodevelopmental effects of long-term therapy. *J Pediatr* 113:58–64.
- Budd MA, Tanaka K, Holmes LB, Efron ML, Crawford ML, Isselbacher KJ. 1967. Isovaleric acidemia: Clinical features of a new genetic defect of leucine metabolism. *NEJ Med* 277:321–327.
- Burke DG, Halpern B, Malegan D, McCairns E, Danks D, Schlesinger P, Wilken B. 1983. Profiles of urinary volatiles from metabolic disorders characterized by unusual odors. *Clin Chem* 29:1834–1838.
- Cohn RM, Yudkoff M, Rothman R, Segal S. 1978. Isovaleric acidemia: Use of glycine therapy in neonates. *NEJ Med* 299:996–999.
- Collins JE, Uempleby AM, Boroujerdi MA, Leonard JV, Sonksen PH. 1987. Effect of insulin on leucine kinetics in maple syrup urine disease. *Pediatr Res* 21:10–13.
- Coude FX, Sweetman L, Nyhan WL. 1979. Inhibition by propionyl-coenzyme A of N-acetylglutamate synthetase in rat liver mitochondria. A possible explanation for hyperammonemia in propionic and methylmalonic acidemia. *J Clin Inv* 64:1544–1551.
- Crane FL, Mii S, Hauge JG, Green DE, Beinert H. 1956. On the mechanism of dehydrogenation of fatty acyl derivatives of coenzyme A. I. The general fatty acyl coenzyme A dehydrogenase. *J Biol Chem* 218:701–706.
- Dorland L, Duran M, Wadman SK, Niederwieser A, Bruinvis L, Ketting D. 1983. Isovalerylglycuronide, a new urinary metabolite in isovaleric acidemia. Identification problems due to rearrangement reactions. *Clinica Chimica Acta* 134:77–83.

- Efron ML. 1967. Isovaleric acidemia. *Am J Disease of Children* 113:74–76.
- Elsas LJ, II, Naglak M. 1988. Acute and chronic-intermittent isovaleric acidemia: Diagnosis and glycine therapy. *Acta Paediatrica Japonica* 30:442–451.
- Engel PC, Massey V. 1971. Green butyryl-coenzyme A dehydrogenase. An enzyme-acyl-coenzyme A complex. *Bioch J* 125: 889–902.
- Ensenauer R, Vockley J, Willard JM, Huey JC, Sass JO, Edland SD, Burton BK, Berry SA, Santer R, Grunert S, Koch HG, Marquardt I, Rinaldo P, Hahn S, Matern D. 2004. A common mutation is associated with a mild, potentially asymptomatic phenotype in patients with isovaleric acidemia diagnosed by newborn screening. *Am J Hum Genet* 75:1136–1142.
- Ensenauer R, He M, Willard JM, Goetzman ES, Corydon TJ, Vandahl BB, Mohsen AW, Isaya G, Vockley J. 2005. Human acyl-CoA dehydrogenase-9 plays a novel role in the mitochondrial beta-oxidation of unsaturated fatty acids. *J Biol Chem* 280:32309–32316.
- Feinstein JA, O'Brien K. 2003. Acute metabolic decompensation in an adult patient with isovaleric acidemia. *S Med J* 96:500–503.
- Fischer AQ, Challa VR, Burton BK, McLean WT. 1981. Cerebellar hemorrhage complicating isovaleric acidemia: A case report. *Neurology* 31:746–748.
- Frerman FE, Kim JJP, Huhta K, McKean MC. 1980. Properties of the general acyl-CoA dehydrogenase from pig liver. *J Biol Chem* 255:2195–2198.
- Fries MH, Rinaldo P, Schmidt-Sommerfeld E, Jurecki E, Packman S. 1996. Isovaleric acidemia: Response to a leucine load after three weeks of supplementation with glycine, L-carnitine, and combined glycine-carnitine therapy. *J Pediatr* 129:449–452.
- Ghisla S, Massey V. 1989. Mechanisms of flavoprotein-catalyzed reactions. *Euro J Biochem* 181:1–17.
- Ghisla S, Engst S, Moll M, Bross P, Strauss AW, Kim JJP. 1992. a,b-Dehydrogenation by acyl-CoA dehydrogenases: Role of functional groups at the active center. *Prog Clin Biol Res—New Dev In Fatty Acid Oxidation* 375:127–142.
- Gibson KM, Burlingame TG, Hogema B, Jakobs C, Schutgens RBH, Millington D, Roe CR, Roe DS, Sweetman L, Steiner RD, Linck L, Pohowalla P, Sacks M, Kiss D, Rinaldo P, Vockley J. 2000. 2-Methylbutyryl-coenzyme A dehydrogenase deficiency: A new inborn error of L-isoleucine metabolism. *Pediatr Res* 47:830–833.
- Gilbert-Barnes E, Barnes LA. 1999. Isovaleric acidemia with promyelocytic myeloproliferative syndrome. *Pediatr Dev Path* 2:286–291.
- Gorelick RJ, Schopfer LM, Ballou DP, Massey V, Thorpe C. 1985. Interflavin oxidation-reduction reactions between pig kidney general acyl-CoA dehydrogenase and electron-transferring flavoprotein. *Biochemistry* 24:6830–6839.
- Harris RA, Joshi M, Jeoung NH. 2004. Mechanisms responsible for regulation of branched-chain amino acid catabolism. *Biochem Biophys Res Commun* 313:391–396.
- Hine DG, Tanaka K. 1984. The identification and the excretion pattern of isovaleryl glucuronide in the urine of patients with isovaleric acidemia. *Pediatr Res* 18:508–512.
- Hine DG, Hack AM, Goodman SI, Tanaka K. 1986. Stable isotope dilution analysis of isovalerylglycine in amniotic fluid and urine and its application for the prenatal diagnosis of isovaleric acidemia. *Pediatr Res* 20:222–226.
- Hyman DB, Tanaka K. 1986. Isovaleryl-CoA dehydrogenase activity in isovaleric acidemia fibroblasts using an improved tritium release assay. *Pediatr Res* 20:59–61.
- Ikeda Y, Tanaka K. 1988. Isovaleryl-CoA dehydrogenase from rat liver. *Methods Enzymol* 166:374–389.
- Ikeda Y, Dabrowski C, Tanaka K. 1983. Separation and properties of five distinct acyl-CoA dehydrogenases from rat liver mitochondria. *J Biol Chem* 258:1066–1076.
- Ikeda Y, Hine DG, Okamura-Ikeda K, Tanaka K. 1985a. Mechanism of action of short-chain, medium-chain, and long-chain acyl-CoA dehydrogenases: Direct evidence for carbanion formation as an intermediate step using enzyme-catalyzed C-2 proton/deuteron exchange in the absence of C-3 exchange. *J Biol Chem* 260:1326–1337.
- Ikeda Y, Keese S, Tanaka K. 1985b. Molecular heterogeneity of variant isovaleryl acyl-CoA dehydrogenase from cultured isovaleric acidemia fibroblasts. *Proc Natl Acad Sci USA* 82:7081–7085.
- Ikeda Y, Keese S, Fenton WA, Tanaka K. 1987. Biosynthesis of four rat liver mitochondrial acyl-CoA dehydrogenases. Import into mitochondria and processing of their precursors in a cell-free system and in cultured cells. *Arch Biochem Biophys* 252:662–674.
- Itoh T, Ito T, Ohba S, Sugiyama N, Mizuguchi K, Yamaguchi S, Kidouchi K. 1996. Effect of carnitine administration on glycine metabolism in patients with isovaleric acidemia: Significance of acetylcarnitine determination to estimate the proper carnitine dose. *Tohoku J Exp Med* 179:101–109.
- Jakobs C, Sweetman L, Nyhan WL. 1984. Hydroxy acid metabolites of branched-chain amino acids in amniotic fluid. *Clinica Chimica Acta* 140:157–166.
- Kahler SG, Sherwood WG, Woolf D, Lawless ST, Zaritsky A, Bonham J, Taylor CJ, Clarke JT, Durie P, Leonard JV. 1994. Pancreatitis in patients with organic acidemias. *J Pediatr* 124:239–243.
- Kelleher JF Jr, Yudkoff M, Hutchinson R, August CS, Cohn RM. 1980. The pancytopenia of isovaleric acidemia. *Pediatrics* 65:1023–1027.
- Kleij W, Van der Kraan M, Huijman J, Van den Heuvel C, Jakobs C. 1995. Prenatal diagnosis of isovaleric acidemia by enzyme and metabolite assay in the first and second trimesters. *Pren Diagn* 15:527–533.
- Krieger I, Tanaka K. 1976. Therapeutic effects of glycine in isovaleric acidemia. *Pediatr Res* 10:25–29.
- Lehnert W. 1981a. 3-Hydroxyisovaleric acid: A new metabolite in isovaleric acidemia. *Clinica Chimica Acta* 113:101–103.
- Lehnert W. 1981b. Excretion of N-isovalerylglycine in isovaleric acidemia. *Clinica Chimica Acta* 116:249–252.
- Lehnert W. 1983. N-Isovaleryllalanine and N-isovalerylsarcosine: Two new minor metabolites in isovaleric acidemia. *Clinica Chimica Acta* 134:207–212.
- Lehnert W, Niederhoff H. 1981. 4-Hydroxyisovaleric acid: A new metabolite in isovaleric acidemia. *Euro J Pediatr* 136:281–283.
- Levy HL, Erickson AM, Lott IT, Kurtz DJ. 1973. Isovaleric acidemia: Results of family study and dietary treatment. *Pediatrics* 52:83–94.
- Loots DT, Erasmus E, Mienie LJ. 2005. Identification of 19 new metabolites by induced by abnormal amino acid conjugation in isovaleric acidemia. *Clin Chem* 51:1510–1512.
- Lott IT, Erickson AM, Levy HL. 1972. Dietary treatment of an infant with isovaleric acidemia. *Pediatrics* 49:616–618.
- Mayatepek E, Kurczynski TW, Hoppel CL. 1991. Long-term L-carnitine treatment in isovaleric acidemia. *Pediatric Neuro* 7:137–140.
- McKean MC, Frerman FE, Mielke DM. 1979. General acyl-CoA dehydrogenase from pig liver. Kinetic and binding studies. *J. Biol. Chem* 254:2730–2735.
- Mehta KC, Zsolway K, Osterhoudt KC, Krantz I, Henretig FM, Kaplan P. 1996. Lessons from the late diagnosis of isovaleric acidemia in a five-year-old boy. *J Pediatr* 129:309–310.
- Mendiola J Jr, Robotham JL, Liehr JG, Williams JC. 1984. Neonatal lethargy due to isovaleric acidemia and hyperammonemia. *Tex Med* 80:52–54.
- Millington DS, Roe CR, Maltby DA, Inoue F. 1987. Endogenous catabolism is the major source of toxic metabolites in isovaleric acidemia. *J Pediatr* 110:56–60.
- Miura R, Nishina Y, Fujii S, Shiga K. 1996. C-13-Nmr Study On the Interaction Of Medium-Chain Acyl-Coa Dehydrogenase With Acetoacetyl-Coa. *J Biochem* 119:512–519.
- Mohsen A-WA, Vockley J. 1995. Identification of the active site catalytic residue in human isovaleryl-CoA dehydrogenase. *Biochemistry* 34:10146–10152.
- Mohsen AW, Anderson BD, Volchenbom SL, Battaile KP, Tiffany K, Roberts D, Kim JJ, Vockley J. 1998. Characterization of molecular defects in isovaleryl-CoA dehydrogenase in patients with isovaleric acidemia. *Biochemistry* 37:10325–10335.
- Nagao M, Tanaka K. 1992. FAD-Dependent regulation of transcription, translation, post-translational processing, and post-processing stability of various mitochondrial Acyl-CoA dehydrogenases and of electron transfer flavoprotein and the site of holoenzyme formation. *J Biol Chem* 267:17925–17932.
- Nasser I, Mohsen AW, Jelezarov I, Vockley J, Macheroux P, Ghisla S. 2004. Thermal unfolding of medium-chain acyl-CoA dehydrogenase and iso(3)valeryl-CoA dehydrogenase: Study of the effect of genetic defects on enzyme stability. *Biochim Biophys Acta* 1690:22–32.
- Nishina Y, Sato K, Hazeckawa I, Shiga K. 1995. Structural modulation of 2-enoyl-CoA bound to reduced acyl-CoA dehydrogenases: A resonance Raman study of a catalytic intermediate. *J Biochem* 117:800–808.
- Parimoo B, Tanaka K. 1993. Structural organization of the human isovaleryl-CoA dehydrogenase gene. *Genomics* 15:582–590.

- Pollitt RJ. 1987. Endogenous catabolism as source of toxic metabolites in isovaleric acidemia. *J Pediatr* 111:477–478.
- Poorthuis BJ, Jille-Vlckova T, Onkenhout W. 1993. Determination of acylcarnitines in urine of patients with inborn errors of metabolism using high-performance liquid chromatography after derivatization with 4'-bromophenacylbromide. *Clinica Chimica Acta* 216:53–61.
- Rhead WJ, Tanaka K. 1980. Demonstration of a specific mitochondrial isovaleryl-CoA dehydrogenase deficiency in fibroblasts from patients with isovaleric acidemia. *Proc Natl Acad Sci USA* 77:580–583.
- Shigematsu Y, Sudo M, Momoi T, Inoue Y, Suzuki Y, Kameyama J. 1982. Changing plasma and urinary organic acid levels in a patient with isovaleric acidemia during an attack. *Pediatr Res* 16:771–775.
- Shigematsu Y, Kikawa Y, Sudo M, Kanaoka H, Fujioka M, Dan M. 1991. Prenatal diagnosis of isovaleric acidemia by fast atom bombardment and tandem mass spectrometry. *Clinica Chimica Acta* 203:369–374.
- Shih VE, Mandell R, Tanaka K. 1973. Diagnosis of isovaleric acidemia in cultured fibroblasts. *Clinica Chimica Acta* 48:437–439.
- Shih VE, Aubry RH, DeGrande G, Gursky SF, Tanaka K. 1984. Maternal isovaleric acidemia. *J Pediatr* 105:77–78.
- Sogut A, Acun C, Aydin K, Tomsac N, Demirel F, Aktuglu C. 2004. Isovaleric acidemia: Cranial CT and MRI findings. *Pediatr Radiol* 34:1.
- Spinty S, Rogozinski H, Lealman GT, Wraith JE. 2002. Second case of a successful pregnancy in maternal isovaleric acidemia. *J Inher Metab Dis* 25:697–698.
- Spirer Z, Swirsky-Fein S, Zakut V, Legum C, Bogair N, Charles R, Gil-Av E. 1975. Acute neonatal isovaleric acidemia. A report of two cases. *Israel J Med Sci* 11:1005–1010.
- Stewart PM, Walsler M. 1980. Failure of the normal ureagenic response to amino acids in organic acid-loaded rats. Proposed mechanism for the hyperammonemia of propionic and methylmalonic acidemia. *J Clin Invest* 66:484–492.
- Sweetman L, Williams JD. 2001. Branched chain organic acidurias. In: Scriver C, Beaudet AL, Sly W, Valle D, editors. *The metabolic and molecular basis of inherited disease*, 8th edition. New York: McGraw-Hill. pp 2125–2164.
- Tajima G, Sakura N, Yofune H, Dwi Bahagia Febriani A, Nishimura Y, Sakamoto A, Ono H, Shigematsu Y, Kobayashi M. 2005. Establishment of a practical enzymatic assay method for determination of isovaleryl-CoA dehydrogenase activity using high-performance liquid chromatography. *Clinica Chimica Acta* 353:193–199.
- Tanaka K. 1990. Isovaleric acidemia: Personal history, clinical survey and study of the molecular basis. *Prog Clin Biol Res* 321:273–290.
- Tanaka K, Isselbacher KJ. 1967. The isolation and identification of N-isovalerylglycine from urine of patients with isovaleric acidemia. *J Biol Chem* 242:2966–2972.
- Tanaka K, Budd MA, Efron ML, Isselbacher KJ. 1966. Isovaleric acidemia: A new genetic defect of leucine metabolism. *Proc Natl Acad Sci USA* 56:236–242.
- Tanaka K, Orr JC, Isselbacher KJ. 1968. Identification of beta-hydroxyisovaleric acid in the urine of a patient with isovaleric acidemia. *Biochimica et Biophysica Acta* 152:638–641.
- Tanaka K, Hine DG, West-Dull A, Lynn TB. 1980. Gas-chromatographic method of analysis for urinary organic acids. I. Retention indices of 155 metabolically important compounds. *Clin Chem* 26:1839–1846.
- Vockley J, Parimoo B, Tanaka K. 1991. Molecular characterization of four different classes of mutations in the isovaleryl-CoA dehydrogenase gene responsible for isovaleric acidemia. *Am J Hum Genet* 40:147–157.
- Vockley J, Nagao M, Parimoo B, Tanaka K. 1992a. The variant human isovaleryl-CoA dehydrogenase gene responsible for type-II isovaleric acidemia determines an RNA splicing error, leading to the deletion of the entire 2nd coding exon and the production of a truncated precursor protein that interacts poorly with mitochondrial import receptors. *J Biol Chem* 267:2494–2501.
- Vockley J, Parimoo B, Nagao M, Tanaka K. 1992b. Identification of the molecular defects responsible for the various genotypes of isovaleric acidemia. *Prog Clin Biol Res* 375:533–540.
- Vockley J, Rogan PK, Anderson BD, Willard J, Seelan RS, Smith DI, Liu WG. 2000. Exon skipping in IVD RNA processing in isovaleric acidemia caused by point mutations in the coding region of the IVD gene. *Am J Hum Gen* 66:356–367.
- Weinberg GL, Laurito CE, Geldner P, Pygon BH, Burton BK. 1997. Malignant ventricular dysrhythmias in a patient with isovaleric acidemia receiving general and local anesthesia for suction lipectomy. *J Clin Anesth* 9:668–670.
- Worthen HG, al Ashwal A, Ozand PT, Garawi S, Rahbeeni Z, al Odaib A, Subramanyam SB, Rashed M. 1994. Comparative frequency and severity of hypoglycemia in selected organic acidemias, branched chain amino acidemia, and disorders of fructose metabolism. *Brain Dev* 16:81–85.
- Yoshida I, Sovik O, Sweetman L, Nyhan WL. 1985. Metabolism of leucine in fibroblasts from patients with deficiencies in each of the major catabolic enzymes: Branched-chain ketoacid dehydrogenase, isovaleryl-CoA dehydrogenase, 3-methylcrotonyl-CoA carboxylase, 3-methylglutaconyl-CoA hydratase, and 3-hydroxy-3-methylglutaryl-CoA lyase. *J Neurogen* 2:413–424.
- Yoshino M, Yoshida I, Yamashita F, Mori M, Uchiyama C, Tatibana M. 1982. Neonatal isovaleric acidemia associated with hyperammonemia. *Adv Exp Med Biol* 153:141–146.
- Yudkoff M, Cohn RM, Puschak R, Rothman R, Segal S. 1978. Glycine therapy in isovaleric acidemia. *J Pediatr* 92:813–817.

[Q1](#): Please check the change made in the reference Ghisla et al., 1992.